

UNDERGRADUATE STUDENT Reference Group

# Executive Summary Fall 2020

The ACHA-NCHA III supports the health of the campus community by fulfilling the academic mission, supporting short- and long-term healthy behaviors, and gaining a current profile of health trends within the campus community.



Copyright 2021

All Rights Reserved

## **Table of Contents**

II. Findings	
A. General Health and Campus Climate	
B. Nutrition, BMI, Physical Activity, and Food Security	
C. Health Care Utilization	
D. Impediments to Academic Performance	
E. Violence, Abusive Relationships, and Personal Safety	
F. Tobacco, Alcohol, and Other Drug Use	
G. Sexual Behavior	
H. Mental Health and Wellbeing	
I. Acute Conditions	
J. Ongoing or Chronic Conditions	
K. Sleep	

ACHA, the nation's principal advocate and leadership organization for college and university health, represents a diverse membership that provides and supports the delivery of health care and prevention and wellness services for the nation's 20 million college students. For more information about the association's programs and services, visit www.acha.org, and www.acha.org/NCHA.

Suggested citation for this document:

American College Health Association. American College Health Association-National College Health Assessment III: Undergraduate Student Reference Group Executive Summary Fall 2020. Silver Spring, MD: American College Health Association; 2020.

## **Introduction and Notes**

The ACHA-National College Health Assessment (ACHA-NCHA) is a national research survey organized by the American College Health Association (ACHA) to assist college health service providers, health educators, counselors, and administrators in collecting data about their students' habits and behaviors on the most prevalent health topics. The ACHA-NCHA now provides the largest known comprehensive data set on the health of college students, providing the college health and higher education fields with a vast spectrum of information on student health.

ACHA initiated the original ACHA-NCHA in 2000 and the instrument was used nationwide through the Spring 2008 data collection period. A revised survey, the ACHA-NCHA-II, was in use from Fall 2008 - Spring 2019 data collection periods. The survey was redesigned again, and data collection with the ACHA-NCHA III began in Fall 2019.

Please note that it is not appropriate to compare trends between versions of the survey. Directly comparing data points between the Original ACHA-NCHA, the ACHA-NCHA II, and the ACHA-NCHA III can lead to an erroneous conclusions and is not recommended.

#### Notes about this report:

1. Missing values have been excluded from analysis and only valid percents are included in this document, unless otherwise noted.

2. The ACHA-NCHA III is programmed differently than earlier versions of the survey. Rather than asking the respondents to answer every question (and offering a "not applicable" option), display logic was used throughout the survey to determine whether, based on their response to an earlier question, the student saw a follow-up question. This makes the valid percents of certain questions impossible to apply to the entire sample, as the denominator used was limited to only the number of students that saw the question. When appropriate, results are also presented using the entire sample as the denominator to show the proportion of the overall sample that experienced a particular issue. These differences in presentation are carefully noted throughout the document and will often explain differences observed between this document and the full data report. Please look carefully at descriptions of the data presented in each table, as well as any footnotes included.

3. *About the use of sex and gender in this report:* Survey responses are reported by sex based on the responses to questions 67A, 67B, and 67C. For the purpose of the ACHA-NCHA report documents, respondents are reported as male or female only when their responses to these three questions are consistent with one another. If gender identity is consistent with sex at birth AND "no" is selected for transgender, then respondents are designated as either *male* or *female*. If respondents select "yes" for transgender OR their sex at birth is not consistent with their gender identity, then they are designated as *non-binary*. A respondent that selects "intersex" for sex at birth, "no" for transgender, and man or woman for gender identity are designated as *male* or *female*. A respondent that selects "intersex" for sex at birth, "yes" for transgender, or selects a gender identity other than man or woman are designated as *non-binary*. A respondent that skips any of the three questions is designated as *unknown*. Totals displayed in this report include non-binary and unknown responses.

For additional information about the survey's development, design, and methodology, email Mary T Hoban, PhD, MCHES, (mhoban@acha.org), Christine Kukich, MS (ckukich@acha.org), or visit www.acha-ncha.org.

This Executive Summary highlights results of the ACHA-NCHA III Fall 2020 survey for Undergraduate Student Reference Group consisting of 8,851 respondents.

The response rate was 13.9%.

## A. General Health and Campus Climate

- 58.1 % of college students surveyed ( 65.6 % male and *very good or excellent.*
- 89.8 % of college students surveyed ( 92.0 % male and *good, very good or excellent*.
- 56.5 % female) described their health as
- 89.7 % female) described their health as

.

.

#### Proportion of college students who reported they agree or strongly agree that:

Percent (%)	Male	Female	Total
I feel that I belong at my college/university	70.1	71.9	70.8
I feel that students' health and well-being is a priority at my college/university	61.1	61.3	60.7
At my college/university, I feel that the campus climate encourages free and open			
discussion of students' health and well-being.	60.9	62.4	61.4
At my college/university, we are a campus where we look out for each other	54.2	57.2	55.7

## B. Nutrition, BMI, Physical Activity, and Food Security

#### **College students reported:**

Percent (%)	Male	Female	Total
Drinking 0 sugar-sweetened beverages (per day), on average, in the last 7 days	29.5	29.8	29.5
Drinking 1 or more sugar-sweetened beverages (per day), on average, in the last 7 days	70.5	70.2	70.5
Drinking energy drinks or shots on 0 of the past 30 days	72.8	79.9	77.8
Drinking energy drinks or shots on 1-4 of the past 30 days	15.6	11.9	13.0
Drinking energy drinks or shots on 5 or more of the past 30 days	11.6	8.2	9.2
Eating 3 or more servings of fruits (per day), on average, in the last 7 days	17.7	18.0	17.8
Eating 3 or more servings of vegetables (per day), on average, in the last 7 days	29.1	30.1	29.7

**Estimated Body Mass Index (BMI):** This figure incorporates reported height and weight to form a general indicator of physical health. Categories defined by The World Health Organization (WHO) 2000, reprinted 2004. Obesity: Preventing and Managing the Global Epidemic. WHO Tech Report Series: 894.

BMI Percent (%)	Male	Female	Total
<18.5 Underweight	5.8	5.2	5.4
18.5-24.9 Healthy Weight	49.9	56.3	54.3
25-29.9 Overweight	28.9	21.1	23.3
30-34.9 Class I Obesity	9.5	9.6	9.7
35-39.9 Class II Obesity	3.8	4.0	4.1
≥40 Class III Obesity	2.0	3.7	3.4
Mean	25.29	25.24	25.30
Median	24.33	23.49	23.73
Std Dev	5.46	6.34	6.18

#### Students meeting the recommended guidelines for physical activity

Based on: US Dept of Health and Human Services. *Physical Activities Guidelines for Americans*, 2nd edition. Washington, DC: US Dept of Health and Human Services; 2018

Definitions:

- Recommendation for aerobic activity: 150 minutes or more of moderate-intensity physical activity per week or 75 minutes of vigorous-intensity physical activity or the equivalent combination
- Recommendation for strength training: 2 or more days a week of moderate or greater intensity activities that involve all major muscle groups
- Active Adults meet the recommendation for strength training AND aerobic activity
- Highly Active Adults meet the recommendation for strength training and TWICE the recommendation for aerobic activity (300 minutes or more of moderate-intensity physical activity per week or 150 minutes of vigorous-intensity physical activity or the equivalent combination)

Percent (%)	Male	Female	Total
Guidelines met for aerobic exercise only	70.2	63.1	64.5
Guidelines met for Active Adults	47.0	35.1	37.8
Guidelines met for Highly Active Adults	37.7	26.7	29.2

#### **Food Security**

Based on responses to the US Household Food Security Survey Module: Six-Item Short Form (2012) from the USDA Economic Research Service.

	Percent (%)	Male	Female	Total
High or marginal food security (score 0-1)		63.8	59.9	60.5
Low food security (score 2-4)		21.6	23.7	23.3
Very low food security (score 5-6)		14.6	16.4	16.2
Any food insecurity (low or very low food security)		36.2	40.1	39.5

## **C. Health Care Utilization**

#### **College students reported:**

Pe	ercent (%)	Male	Female	Total
Receiving psychological or mental health services within the last 12 months		18.6	29.8	27.7
*The services were provided by:				
My current campus health and/or counseling center		41.3	38.7	39.9
A mental health provider in the local community near my campus		19.8	22.6	22.3
A mental health provider in my home town		50.1	53.7	52.6
A mental health provider not described above		11.2	8.1	8.7
*Only students who reported receiving care in the last 12 months were asked these questions	s			
	s ercent (%)	Male	Female	Total
		<b>Male</b> 55.1	<b>Female</b> 71.6	<b>Total</b> 67.4
Pe				
Visiting a medical provider within the last 12 months				
Visiting a medical provider within the last 12 months         *The services were provided by:		55.1	71.6	67.4
Visiting a medical provider within the last 12 months         *The services were provided by:         My current campus health center		55.1 23.7	22.7	67.4

\*Only students who reported receiving care in the last 12 months were asked these questions

#### **College students reported:**

Percent (%)	Male	Female	Total
Flu vaccine within the last 12 month	51.1	56.5	55.0
Not starting the HPV vaccine series	18.7	16.0	16.5
Starting, but not completing HPV vaccine series	4.0	6.0	5.5
Completing HPV vaccine series	44.6	56.9	53.6
Not knowing their HPV vaccine status	32.7	21.1	24.4
Ever having a GYN visit or exam (females only)		47.9	
Having a dental exam in the last 12 months	68.7	73.2	71.6
Being tested for HIV within the last 12 months	7.2	10.8	9.9
Being tested for HIV more than 12 months ago	6.4	8.1	7.8
Wearing sunscreen usually or always when outdoors	16.6	37.2	31.4
Spending time outdoors with the intention of tanning at least once in the last 12 months	39.6	65.3	57.6

### **D. Impediments to Academic Performance**

Respondents are asked in numerous places throughout the survey about issues that might have negatively impacted their academic performance within the last 12 months. This is defined as negatively impacting their performance in a class or delaying progress towards their degree. Both types of negative impacts are represented in the figures below. Please refer to the corresponding Data Report for specific figures on each type of impact. Figures in the left columns use all students in the sample as the denominator. Figures in the right columns use only the students that experienced that issue (e.g. students who used cannabis, reported a problem or challenge with finances, or experienced a particular health issue) in the denominator. (items are listed in the order in which they appear in the survey)

			tively imp nic perfor				tively impរ mic perforı	
		among <u>all students in the</u> sample				0	<u>only studer</u> rienced the	
	Percent (%)	Male	Female	Total	1	Male	Female	Total
Alcohol use		2.0	1.4	1.5		3.1	1.9	2.2
Cannabis/marijuana use		1.7	1.1	1.4		5.8	3.9	4.0

#### Problems or challenges in the last 12 months

Troblems of chanenges in the last 12 months			
Career	11.2	12.0	12.1
Finances	15.0	17.8	17.4
Procrastination	49.7	49.1	49.8
Faculty	7.7	6.9	7.1
Family	7.7	13.6	12.4
Intimate Relationships	8.8	10.2	10.0
Roommate/housemate	5.0	6.1	5.9
Peers	3.6	4.0	3.9
Personal appearance	3.8	6.2	6.0
Health of someone close to me	6.7	10.9	10.0
Death of a family member, friend, or someone close to			
me	7.0	10.6	9.7
Bullying	1.0	1.7	1.6
Cyberbullying	0.7	1.0	0.9
Hazing	0.3	0.2	0.3
Microaggression	1.4	2.0	2.0
Sexual Harassment	0.5	2.2	1.8
Discrimination	1.1	1.7	1.7

38.3	36.3	37.2
38.9	36.1	37.1
67.0	63.7	65.1
62.5	56.9	57.7
29.5	33.5	33.0
26.0	28.3	28.0
25.3	21.9	23.0
18.3	16.1	16.7
11.3	11.2	11.9
23.9	27.9	27.5
35.6	42.3	41.0
24.7	27.7	27.0
28.3	25.4	25.3
38.1	43.8	42.1
15.3	14.2	14.6
20.8	20.1	20.7
18.1	18.0	18.6

4.6

	Negatively impacted academic performance among <u>all students in the</u> <u>sample</u>				acader among	tively impa mic perform only studen ienced the	formance dents that	
Percent (%)	Male	<u>sample</u> Female	Total	1	Male	Female	Total	
	marc	remate	Total	L	Wate	remare	Total	
Acute Diagnoses in the last 12 months Bronchitis	0.6	1.4	1.2	Г	35.0	42.9	42.3	
Chlamydia	0.0	0.2	1.2 0.1	_	11.1	13.3	42.3	
Chicken Pox (Varicella)								
	0.0	0.0	0.0		12.5	7.7	9.5	
Cold/Virus or other respiratory illness	5.8	11.4			26.8	34.9	33.5	
Concussion	0.8	2.3	2.0	_	39.6	65.0	61.6	
Gonorrhea	0.1	0.0	0.0		20.0	12.5	14.8	
Flu (influenza or flu-like illness)	2.6	4.0	3.8		32.8	46.7	44.0	
Mumps	0.0	0.0	0.0		11.1	0.0	3.8	
Mononucleosis (mono)	0.4	0.9	0.8		40.9	51.9	51.1	
Orthopedic injury	1.5	2.9	2.5		22.1	29.0	27.5	
Pelvic Inflammatory Disease	0.0	0.1	0.1		12.5	26.9	22.9	
Pneumonia	0.6	0.7	0.7		39.4	45.0	44.3	
Shingles	0.2	0.1	0.1		28.6	20.0	22.5	
Stomach or GI virus or bug, food poisoning or gastritis	1.9	3.6	3.2		31.4	38.7	37.5	
Urinary tract infection	0.0	2.4	1.8		5.9	17.5	17.3	
Percent (%)	Male	Female	Total		Male	Female	Total	
Any ongoing or chronic medical conditions diagnosed or treated in the last 12 months	18.4	30.0	27.7		27.6	38.5	36.9	
Other impediments to academic performance								
Assault (physical)	0.2	0.9	0.8		8.6	30.2	25.7	
Assault (sexual)	0.6	2.3	2.0		25.0	29.6	30.6	
Allergies	2.3	2.8	2.7	[	6.7	6.8	6.8	
Anxiety	23.4	38.5	35.2		46.9	54.3	53.6	
ADHD or ADD	9.8	9.8	10.3		60.1	67.3	66.1	
Concussion or TBI	1.3	2.4	2.1		37.5	51.4	48.7	

(	Other	imped	liments	to	academic	performa	ice
	· · · · · ·			•••		P	

Assault (physical)	0.2	0.9	0.8
Assault (sexual)	0.6	2.3	2.0
Allergies	2.3	2.8	2.7
Anxiety	23.4	38.5	35.2
ADHD or ADD	9.8	9.8	10.3
Concussion or TBI	1.3	2.4	2.1
Depression	19.0	27.8	26.4
Eating disorder/problem	1.6	4.4	3.9
Headaches/migraines	6.5	16.9	14.3
Influenza or influenza-like illness (the flu)	2.6	4.4	4.0
Injury	1.8	2.3	2.2
PMS	0.1	14.1	10.5
PTSD	1.8	3.7	3.4
Short-term illness	2.0	4.7	4.0
Upper respiratory illness	3.4	7.2	6.3
Sleep difficulties	21.2	26.5	25.7
Stress	32.1	49.1	45.2

8.6	30.2	25.7
25.0	29.6	30.6
6.7	6.8	6.8
46.9	54.3	53.6
60.1	67.3	66.1
37.5	51.4	48.7
53.3	60.0	59.6
24.2	27.7	28.4
23.7	36.3	34.2
28.5	38.2	36.3
16.1	21.0	19.8
13.3	26.4	26.7
49.4	42.0	44.6
19.9	31.7	29.5
19.4	24.1	23.7
46.1	46.7	47.3
46.9	57.3	55.6

## E. Violence, Abusive Relationships, and Personal Safety

### Within the last 12 months, college students reported experiencing:

Within the last 12 months, college students reported experiencing:			-
Percent (%)	Male	Female	Total
A physical fight	4.3	1.9	2.7
A physical assault (not sexual assault)	1.8	2.1	2.1
A verbal threat	14.7	10.2	11.8
Sexual touching without their consent	2.0	7.0	5.8
Sexual penetration attempt without their consent	0.8	3.1	2.6
Sexual penetration without their consent	0.7	2.2	1.8
Being a victim of stalking	1.6	5.1	4.1
A partner called me names, insulted me, or put me down to make me feel bad	9.2	12.2	11.3
A partner often insisted on knowing who I was with and where I was or tried to limit my			
contact with family or friends	5.9	6.6	6.5
A partner pushed, grabbed, shoved, slapped, kicked, bit, choked or hit me without my			
consent	2.8	3.1	3.0
A partner forced me into unwanted sexual contact by holding me down or hurting me in			
some way	1.3	2.3	2.0
A partner pressured me into unwanted sexual contact by threatening me, coercing me, or			
using alcohol or other drugs	1.6	3.9	3.3

College students reported feeling *very safe* :

College students reported feeling very saje :			
Percent (%)	Male	Female	Total
On their campus (daytime)	83.3	74.7	76.6
On their campus (nighttime)	49.5	19.0	27.1
In the community surrounding their campus (daytime)	50.9	41.7	43.8
In the community surrounding their campus (nighttime)	26.4	9.7	14.1

## F. Tobacco, Alcohol, and Other Drug Use

	Ever Used			*Used in the last 3 months			
Percent (%)	Male	Female	Total	Male	Female	Total	
Tobacco or nicotine delivery products (cigarettes, e-							
cigarettes, Juul or other vape products, water pipe or							
hookah, chewing tobacco, cigars, etc.)	36.4	31.6	32.8	26.5	21.1	22.6	
Alcoholic beverages (beer, wine, liquor, etc.)	66.1	69.9	68.7	59.4	63.5	62.1	
Cannabis (marijuana, weed, hash, edibles, vaped							
cannabis, etc.) [Please report nonmedical use only.]	35.4	35.7	35.9	24.2	23.6	24.0	
Cocaine (coke, crack, etc.)	6.0	3.9	4.4	2.4	1.1	1.4	
Prescription stimulants (Ritalin, Concerta, Dexedrine,							
Adderall, diet pills, etc.) [Please report nonmedical							
use only.]	9.5	6.2	7.2	3.9	2.7	3.0	
Methamphetamine (speed, crystal meth, ice, etc.)	0.7	0.6	0.7	0.1	0.1	0.1	
Inhalants (poppers, nitrous, glue, gas, paint thinner,							
etc.)	2.6	1.5	1.9	0.9	0.5	0.7	
Sedatives or Sleeping Pills (Valium, Ativan, Xanax,							
Klonopin, Librium, Rohypnol, GHB, etc.) [Please							
report nonmedical use only.]	4.9	3.7	4.1	1.7	1.5	1.6	
Hallucinogens (Ecstasy, MDMA, Molly, LSD, acid,							
mushrooms, PCP, Special K, etc.)	10.3	6.2	7.5	4.4	2.4	3.1	
Heroin	0.5	0.3	0.4	0.1	0.0	0.0	
Prescription opioids (morphine, codeine, fentanyl,							
oxycodone [OxyContin, Percocet], hydrocodone							
[Vicodin], methadone, buprenorphine [Suboxone],							
etc.) [Please report nonmedical use only.]	3.9	2.3	2.8	1.0	0.5	0.6	

\*These figures use all students in the sample as the denominator, rather than just those students who reported lifetime use.

#### Substance Specific Involvement Scores (SSIS) from the ASSIST

	*Moderate risk use of the substance			*High risk use of the substance			
Percent (%)	Male	Female	Total	Male	Female	Total	
Tobacco or nicotine delivery products	19.6	14.2	15.6	1.1	1.0	1.0	
Alcoholic beverages	9.8	9.3	9.4	1.0	0.6	0.7	
Cannabis (nonmedical use)	16.3	15.1	15.6	0.7	0.6	0.7	
Cocaine	1.4	0.7	0.8	0.0	0.0	0.0	
Prescription stimulants (nonmedical use)	1.6	1.5	1.5	0.1	0.0	0.0	
Methamphetamine	0.2	0.2	0.2	0.0	0.0	0.0	
Inhalants	0.4	0.3	0.3	0.0	0.0	0.0	
Sedatives or Sleeping Pills (nonmedical use)	1.2	1.0	1.1	0.0	0.0	0.0	
Hallucinogens	2.4	1.3	1.7	0.0	0.0	0.0	
Heroin	0.1	0.1	0.1	0.0	0.0	0.0	
Prescription opioids (nonmedical use)	0.7	0.4	0.5	 0.0	0.0	0.0	

\*These figures use all students in the sample as the denominator, rather than just those students who reported lifetime use.

Proportion of students (overall sample) who report misusing prescription medications (taking without a prescription, or taking more medication or more often than prescribed) in the past 3 months:

	Percent (%)	Male	Female	Total
Prescription stimulants		2.9	2.2	2.4
Prescription sedatives or sleeping pills		1.3	1.2	1.2
Prescription opioids		0.7	0.3	0.4

#### \*Tobacco or nicotine delivery products used in the last 3 months

Percent (%)	Male	Female	Total
Cigarettes	10.0	5.9	7.2
E-cigarettes or other vape products (for example: Juul,			
etc.)	19.5	16.9	17.5
Water pipe or hookah	2.9	2.0	2.2
Chewing or smokeless tobacco	5.5	0.5	1.9
Cigars or little cigars	8.0	1.5	3.3
Other	0.9	0.2	0.4

\*These figures use all students in the sample as the denominator, rather than just those students who reported tobacco or nicotine delivery product use in the last 3 months.

#### **Students in Recovery**

■ 1.6 % of college students surveyed ( 2.0 % male and from alcohol or other drug use.

#### When, if ever, was the last time you:

	Drank Alcohol				*Used Cannabis/Marijuana			
Percent (%)	Male	Female	Total		Male	Female	Ťotal	
Never	27.2	23.6	24.7		59.7	59.3	59.1	
Within the last 2 weeks	45.7	46.5	46.0		14.9	14.0	14.5	
More than 2 weeks ago but within the last 30 days	7.7	10.4	9.8		3.7	3.8	3.7	
More than 30 days ago but within the last 3 months	7.2	8.5	8.1		5.6	4.6	5.0	
More than 3 months ago but within the last 12 months	6.4	6.9	6.9		6.0	7.5	7.1	
More than 12 months ago	5.8	4.1	4.6		10.2	10.7	10.6	

1.2

% female) indicated they were in recovery

\*Students were instructed to *include* medical and non-medical use of cannabis.

#### Driving under the influence

- 10.5 % of college students reported driving after having *any alcohol* in the last 30 days.\*
  - \*Only students who reported driving in the last 30 days and drinking alcohol in the last 30 days were asked this question.
- 32.7 % of college students reported driving within 6 hours of using cannabis/marijuana in the last 30 days.\* \*Only students who reported driving in the last 30 days and using cannabis in the last 30 days were asked this question.

**Estimated Blood Alcohol Concentration** (or eBAC) of college students. Due to the improbability of a student surviving a drinking episode resulting in an extremely high eBAC, all students with an eBAC of 0.50 or higher are also omitted from these eBAC figures. eBAC is an estimated figure based on the reported number of drinks consumed during the last time they drank alcohol in a social setting, their approximate time of consumption, sex, weight, and the average rate of ethanol metabolism. Only students who reported drinking alcohol within the last 3 months answered these questions.

Estimated BAC	Percent (%)	Male	Female	Total
<.08		78.9	79.4	79.5
<.10		84.1	85.3	85.1
Mean		0.05	0.05	0.05
Median		0.02	0.03	0.02
Std Dev		0.06	0.06	0.06

#### \*Reported number of drinks consumed the last time students drank alcohol in a social setting.

Number of drinks	Percent (%)	Male	Female	Total
4 or fewer		63.0	78.9	75.1
5		8.5	8.8	8.7
6		7.2	4.9	5.4
7 or more		21.2	7.4	10.8
Mean		4.5	3.1	3.4
Median		3.0	2.0	3.0
Std Dev		4.5	2.5	3.2

\*Only students who reported drinking alcohol in the last three months were asked this question.

#### Reported number of times college students consumed five or more drinks in a sitting within the last two weeks:

		ng <u>all</u> stu surveyed		drinking	those who g alcohol wi st two weel	ithin the
Percent (%)	Male	Female	Total	Male	Female	Total
Did not drink alcohol in the last two weeks (includes						
non-drinkers)	54.4	53.5	54.1			
None	19.3	23.8	22.6	42.3	51.2	49.2
1-2 times	17.5	17.1	16.9	38.3	36.8	36.9
3-5 times	6.9	4.7	5.2	15.0	10.1	11.4
6 or more times	2.0	0.8	1.1	4.3	1.8	2.5

\*Only students who reported drinking alcohol in the last two weeks were asked this question.

#### \*College students who drank alcohol reported experiencing the following in the last 12 months when drinking alcohol:

Percent (%)	Male	Female	Total
Did something I later regretted	16.9	17.9	17.6
Blackout (forgot where I was or what I did for a large period of time and cannot			
remember, even when someone reminds me)	12.3	11.1	11.3
Brownout (forgot where I was or what I did for short periods of time, but can			
remember once someone reminds me)	20.4	22.1	21.5
Got in trouble with the police	1.5	0.9	1.0
Got in trouble with college/university authorities	1.0	0.7	0.8
Someone had sex with me without my consent	0.5	2.2	1.8
Had sex with someone without their consent	0.3	0.2	0.2
Had unprotected sex	12.0	13.5	13.0
Physically injured myself	7.1	6.9	7.1
Physically injured another person	1.1	0.6	0.7
Seriously considered suicide	3.4	2.5	2.9
Needed medical help	1.0	0.9	1.0
Reported two or more of the above	23.4	24.1	23.9

\*Only students who reported drinking alcohol in the last 12 months were asked these questions.

## **G. Sexual Behavior**

When, if ever, was the last time you had:		Oral sex	
Percent (%)	Male	Female	Total
Never	39.5	37.2	37.9
Within the last 2 weeks	26.7	32.0	30.4
More than 2 weeks ago but within the last 30 days	5.8	7.6	7.1
More than 30 days ago but within the last 3 months	6.9	7.4	7.2
More than 3 months ago but within the last 12 months	11.2	8.9	9.6
More than 12 months ago	9.8	6.9	7.8

Vaginal intercourse					
Male	Female	Total			
46.5	39.1	41.7			
26.1	35.8	32.6			
4.1	6.2	5.6			
5.4	6.0	5.8			
9.1	7.1	7.6			
8.8	5.8	6.7			

	Ana	Anal intercourse		
Percent (%)	Male	Female	Total	
Never	81.9	83.7	83.1	
Within the last 2 weeks	3.8	1.9	2.5	
More than 2 weeks ago but within the last 30 days	1.1	1.2	1.2	
More than 30 days ago but within the last 3 months	2.2	1.9	2.0	
More than 3 months ago but within the last 12 months	3.4	3.8	3.7	
More than 12 months ago	7.7	7.5	7.5	

## \*College students who reported having oral sex, or vaginal or anal intercourse within the last 12 months reported having the following number of sexual partners:

	Percent (%)	Male	Female	Total
None		0.2	0.2	0.2
1		64.5	68.8	67.7
2		14.9	13.0	13.5
3		8.3	7.5	7.7
4 or more		12.2	10.6	11.0
Mean		2.2	1.8	1.9
Median		1.0	1.0	1.0
Std Dev		3.5	2.0	2.4

\*Only students who reported having oral sex, or vaginal or anal intercourse in the last 12 months were asked this question.

## College students who reported having oral sex, or vaginal or anal intercourse within the last 30 days who reported using a condom or another protective barrier *most of the time* or *always*:

Percent (%)	Male	Female	Total
Oral sex	6.1	5.6	5.8
Vaginal intercourse	44.9	40.7	41.8
Anal intercourse	25.0	22.6	25.3

\*Only students who reported having oral sex, or vaginal or anal intercourse in the last 30 days were asked these questions.

## College students who reported having vaginal intercourse (penis in vagina) within the last 12 months were asked if they or their partner used any method to prevent pregnancy the <u>last time</u> they had vaginal intercourse:

	Percent (%)	Male	Female	Total
Yes, used a method of contraception		85.3	85.8	85.4
No, did not want to prevent pregnancy		1.9	1.9	1.9
No, did not use any method		11.2	11.3	11.4
Don't know		1.6	1.0	1.2

\*Only students who reported having oral sex, or vaginal or anal intercourse in the last 12 months were asked this question.

#### \*Those students who reported using a contraceptive use the last time they had vaginal intercourse, reported they (or their partner) used the following methods: .

Percent (%)	Male	Female	Total
Birth control pills (monthly or extended cycle)	54.4	52.2	52.6
Birth control shots	2.9	2.1	2.4
Birth control implants	8.4	8.1	8.1
Birth control patch	0.8	0.8	0.8
The ring	1.4	2.1	1.9
Emergency contraception ("morning after pill" or "Plan B")	4.9	5.1	5.1
Intrauterine device	11.5	12.1	12.0
Male (external) condom	58.6	49.7	51.9
Female (internal) condom	0.1	0.1	0.2
Diaphragm or cervical cap	0.0	0.1	0.1
Contraceptive sponge	0.1	0.1	0.1
Withdrawal	22.9	28.1	26.9
Fertility awareness (calendar, mucous, basal body temperature)	4.0	5.9	5.4
Sterilization (hysterectomy, tubes tied, vasectomy)	1.3	1.5	1.5
Other method	0.9	1.0	1.0
Male condom use plus another method	40.5	38.0	38.8
Any two or more methods (excluding male condoms)	25.5	27.0	26.7

\*Only students who reported they or their partner used a method the last time they had vaginal intercourse were asked these questions.

College students who reported having vaginal intercourse (penis in vagina) within the last 12 months were asked if they or their partner used emergency contraception ("morning after pill" or "Plan B") in the last 12 months: 19.1 %).

Yes (male: 16.2 %; female:

College students who reported having vaginal intercourse (penis in vagina) within the last 12 months were asked if they experienced an unintentional pregnancy or got someone pregnant within the last 12 months: Yes (male: 0.6 %; female: 1.4 %).

## H. Mental Health and Wellbeing

#### Kessler 6 (K6) Non-Specific Psychological Distress Score (Range is 0-24)\*

	Percent (%)	Male	Female	Total
No or low psychological distress (0-12)		81.8	73.6	74.9
Serious psychological distress (13-24)		18.2	26.4	25.1

\*ACHA discontinued the practice of reporting moderate psychological distress in fall 2023 due to lack of consensus on an appropriate cut off score. Instead, we recommend reporting only students who screen positive (or negative) for serious psychological distress as indicated above. All reference group reports were updated to reflect this change.

Mean	7.72	9.25	8.97
Median	7.00	9.00	8.00
Std Dev	5.25	5.38	5.44

#### UCLA Loneliness Scale (ULS3) Score (Range is 3-9)

	Percent (%)	Male	Female	Total
Negative for loneliness (3-5)		49.7	46.6	46.7
Positive for loneliness (6-9)		50.3	53.4	53.3
Mean		5.49	5.61	5.61
Median		6.00	6.00	6.00
Std Dev		1.89	1.84	1.87

#### **Diener Flourishing Scale – Psychological Well-Being (PWB) Score (Range is 8-56)** (higher scores reflect a higher level of psychological well-being)

	Male	Female	Total
Mean	44.40	45.20	44.73
Median	46.00	47.00	47.00
Std Dev	9.00	8.12	8.56

#### The Connor-Davison Resilience Scale (CD-RISC2) Score (Range is 0-8)

(higher scores reflect greater resilience)

	Male	Female	Total
Mean	6.18	5.84	5.91
Median	6.00	6.00	6.00
Std Dev	1.45	1.46	1.47

#### Self injury

■ 10.7 % of college students surveyed ( 6.4 % male and 11.3 % female) indicated they had intentionally cut, burned, bruised, or otherwise injured themselves within the last 12 months.

## Within the last 12 months, have you had problems or challenges with any of the following:

Male	Female	Total
49.5	51.8	51.9
29.3	33.2	32.8
38.9	49.9	47.2
74.9	77.7	77.3
12.5	12.3	12.6
26.4	40.9	37.9
34.1	36.4	35.9
19.7	28.1	26.0
19.8	25.1	23.9
34.3	56.0	50.8
28.3	39.3	36.8
19.8	25.2	23.9
4.2	6.3	5.9
2.3	3.8	3.6
0.9	0.5	0.6
9.0	14.3	14.0
2.3	11.2	9.0
6.4	9.4	9.2
	49.5 29.3 38.9 74.9 12.5 26.4 34.1 19.7 19.8 34.3 28.3 19.8 4.2 2.3 0.9 9.0 2.3	49.5         51.8           29.3         33.2           38.9         49.9           74.9         77.7           12.5         12.3           26.4         40.9           34.1         36.4           19.7         28.1           19.8         25.1           34.3         56.0           28.3         39.3           19.8         25.2           4.2         6.3           2.3         3.8           0.9         0.5           9.0         14.3           2.3         11.2           6.4         9.4

#### \*Of those reporting this issue, it caused moderate or high distress

ľ	iigh distres	S
Male	Female	Total
82.4	89.9	88.2
68.9	74.5	73.4
74.1	78.7	77.7
60.2	71.0	68.5
56.8	50.9	51.8
59.1	69.3	67.5
62.2	63.4	63.4
44.9	59.3	56.8
41.3	46.1	45.7
49.7	66.1	63.7
58.6	69.5	67.5
68.5	76.9	75.2
42.3	56.2	53.8
35.8	50.4	49.7
42.9	65.6	59.6
25.5	41.1	39.7
35.8	56.0	55.7
39.7	57.4	54.6

\*Only students who reported a problem or challenge in the last 12 months were asked about level of distress.

Students reporting none of the above	9.5	6.3	7.1
Students reporting only one of the above	12.3	6.8	8.1
Students reporting 2 of the above	13.5	9.8	10.5
Students reporting 3 or more of the above	64.7	77.1	74.3

	Percent (%)	Male	Female	Total
Negative suicidal screening (3-6)		76.3	74.0	73.2
Positive suicidal screening (7-18)		23.7	26.0	26.8
Mean		5.05	5.23	5.30
Median		4.00	4.00	4.00
Std Dev		2.91	3.08	3.13

#### Suicide attempt

■ 2.4 % of college students surveyed ( attempted suicide within the last 12 months.

ed ( 2.8 % male and ths

2.1 % female) indicated they had

### Within the last 30 days, how would you rate the overall level of stress experienced:

	Percent (%)	Male	Female	Total
No stress		2.5	0.7	1.2
Low		28.3	12.8	16.6
Moderate		47.6	50.0	49.2
High		21.6	36.5	33.1

### **I. Acute Conditions**

College students reported being diagnosed by a healthcare professional within the last 12 months with:

Percent (%)	Male	Female	Total
Bronchitis	1.9	3.4	3.1
Chlamydia	0.8	1.4	1.2
Chicken Pox (Varicella)	0.4	0.2	0.2
Cold/virus or other respiratory illness (for example:			
sinus infection, ear infection, strep throat, tonsillitis,			
pharyngitis, or laryngitis)	22.7	33.6	30.8
Concussion	2.1	3.7	3.3
Gonorrhea	0.4	0.3	0.3
Flu (influenza) or flu-like illness	8.1	9.0	8.9
Mumps	0.4	0.3	0.3
Mononucleosis (mono)	1.0	1.8	1.6
Orthopedic injury (for example: broken bone, fracture,			
sprain, bursitis, tendinitis, or ligament injury)	7.2	10.3	9.5
Pelvic Inflammatory Disease	0.4	0.4	0.4
Pneumonia	1.6	1.8	1.7
Shingles	0.6	0.4	0.5
Stomach or GI virus or bug, food poisoning or gastritis	6.3	9.9	9.0
Urinary tract infection	1.0	14.3	10.7

## J. Ongoing or Chronic Conditions

The questions for the *ongoing or chronic conditions* are presented differently in this report than the order they appear in the survey. In the survey, all items appear in a single list, ordered alphabetically. In this report, the conditions are presented in groups to ease burden on the reader. The findings are divided into mental health conditions, STIs and other chronic infections, and other ongoing or chronic conditions in this report.

	College students reported <u>ever</u> being diagnosed with the following:				iose rep hea	oorting con lthcare or I ional withir	tact with MH
Mental Health Percent (%)	with Male	the follov Female	ving: Total	l i	months Male Female Tot		
ADD/ADHD - Attention Deficit/Hyperactivity	Whate	remare	Total	1	viaic	remare	10141
Disorder	13.0	9.3	10.6		55.4	64.9	62.1
Alcohol or Other Drug-Related Abuse or Addiction	1.6	0.9	1.2		30.6	42.1	37.3
Anxiety (for example: Generalized Anxiety, Social							
Anxiety, Panic Disorder, Specific Phobia)	17.0	35.0	30.9		63.7	76.2	74.3
Autism Spectrum	2.5	0.8	1.5		32.7	52.2	44.4
Bipolar and Related Conditions (for example: Bipolar							
I, II, Hypomanic Episode)	1.7	2.3	2.3		71.8	75.9	75.2
Borderline Personality Disorder (BPD), Avoidant Personality, Dependent Personality, or another							
personality disorder	0.8	1.2	1.2		38.9	63.4	59.6
Depression (for example: Major depression, persistent depressive disorder, disruptive mood disorder)	13.6	25.3	23.0		69.1	75.1	74.4
Eating Disorders (for example: Anorexia Nervosa,							
Bulimia Nervosa, Binge-Eating)	1.2	5.6	4.6		25.0	54.2	52.0
Gambling Disorder	0.1	0.1	0.1 5.7		0.0	0.0 59.7	0.0
Insomnia	3.0	6.1	5.7		54.5	59.7	59.7
Obsessive-Compulsive and Related Conditions (for example: OCD, Body Dysmorphia, Hoarding, Trichotillomania and other body-focused repetitive behavior disorders)	2.9	6.4	5.7		58.2	60.4	59.8
PTSD (Posttraumatic Stress Disorder), Acute Stress Disorder, Adjustment Disorder, or another trauma- or stressor- related condition	2.7	6.5	5.9		50.8	67.8	65.9
Schizophrenia and Other Psychotic Conditions (for example: Schizophrenia, Schizoaffective Disorder, Schizophreniform Disorder, Delusional Disorder)	0.4	0.3	0.4		60.0	33.3	50.0
Tourette's or other neurodevelopmental condition not							
already listed	0.8	0.4	0.5		31.6	31.8	31.0
Traumatic brain injury (TBI)	0.9	1.3	1.2		33.3	42.9	39.4

\*Only students who reported ever being diagnosed were asked about contact with a healthcare or mental health professional within the last 12 months.

Percent (%)	Male	Female	Total
Students reporting none of the above	68.8	56.6	59.1
Students reporting only one of the above	15.2	14.6	14.7
Students reporting both Depression and Anxiety	10.1	22.1	19.6
Students reporting any two or more of the above			
(excluding the combination of Depression and			
Anxiety)	5.8	6.7	6.6

\*Of those ever diagnosed,

	College students reported <u>ever</u> being diagnosed				had cont or MH	ose ever dia tact with he professiona	althcare l within
STI's/Other chronic infections	with	the follow	ving:		la	st 12 mont	hs
Percent (%)	Male	Female	Total		Male	Female	Total
Genital herpes	0.4	0.9	0.8		33.3	37.5	36.4
Hepatitis B or C	0.4	0.3	0.3		20.0	31.3	25.9
HIV or AIDS	0.2	0.1	0.1		60.0	20.0	40.0
Human papillomavirus (HPV) or genital warts	0.5	1.1	1.0		30.0	43.5	42.2

\*Only students who reported ever being diagnosed were asked about contact with a healthcare or mental health professional within the last 12 months.

					Of the	se ever ula	gne
	College students reported		]	had cont	tact with he	ealt	
	<u>ever</u> k	<u>ever</u> being diagnosed			or MH professional v		ıl w
Other Chronic /Ongoing Medical Conditions	with	with the following:			la	st 12 mont	hs
Percent (%)	Male	Female	Total		Male	Female	]
Acne	25.4	30.0	28.7		33.3	47.2	
Allergies - food allergy	7.9	13.9	12.4		26.8	36.0	
Allergies - animals/pets	11.8	14.4	13.8		24.1	31.6	
Allergies - environmental (for example: pollen, grass,							
dust, mold)	28.8	32.8	31.8		22.5	33.5	
Asthma	16.2	17.6	17.4		22.0	41.2	
Cancer	0.8	0.9			47.1	59.6	
Celiac disease	0.4	1.3	1.1		33.3	48.1	
Chronic pain (for example: back or joint pain, arthritis	,						
nerve pain)	4.7	7.9			63.6	70.2	
Diabetes or pre-diabetes/insulin resistance	2.1	2.4			68.8	63.4	
Endometriosis	0.1	2.5	1.9		0.0	68.0	
Gastroesophageal Reflux Disease (GERD) or acid							
reflux	3.4	6.3	5.5		44.9	54.3	
Heart & vascular disorders (for example: atrial							
fibrillation or other cardiac arrhythmia, mitral valve							
prolapse or other valvular heart disease, congenital							
heart condition)	1.8	-			50.0	54.8	
High blood pressure (hypertension)	3.9	2.0	2.6		61.8	64.3	
High cholesterol (hyperlipidemia)	2.3	3.0	2.8		62.3	54.3	
Irritable bowel syndrome (spastic colon or spastic							
bowel)	2.1	5.6	4.7		41.7	58.2	
Migraine headaches	7.2	15.4	13.4		35.8	55.1	
Polycystic Ovarian Syndrome (PCOS)	0.1	4.1	3.1		50.0	69.6	
Sleep Apnea	1.8	1.4	1.6		59.5	46.0	
Thyroid condition or disorder	1.1	3.7	3.1		65.4	80.4	
Urinary system disorder (for example: bladder or							
kidney disease, urinary reflux, interstitial cystitis)	0.4	1.5	1.2		30.0	47.8	
*Only students who reported over being diagnosed were asked about	t contact wi	th a healthco	re or mento	1 health	professione	l within the la	ct 17

#### \*Of those ever diagnosed, althcare al within

Total 43.3

34.1 29.5 30.5 35.9 55.7 45.7 68.8 64.0 65.5 51.8 52.7 62.8 55.6 57.0 51.8 69.4 52.2 78.2 46.7

\*Only students who reported ever being diagnosed were asked about contact with a healthcare or mental health professional within the last 12 months.

Students who reported being diagnosed with diabetes or pre-diabetes/insulin resistance, indicated they had:

Pei	cent (%)	Male	Female	Total
Type I Diabetes		31.8	22.5	23.9
Type II Diabetes		21.4	17.1	17.4
Pre-diabetes or insulin resistance		58.5	58.3	59.6
Gestational Diabetes		5.0	11.9	9.8

## K. Sleep

Reported amount of time to usually fall asleep at night (sleep onset latency):

Percent (%)	Male	Female	Total
Less than 15 minutes	41.9	36.8	37.8
16 to 30 minutes	27.5	27.3	27.3
31 minutes or more	30.6	35.9	34.9

## Over the last 2 weeks, students reported the following average amount of sleep (excluding naps):

	On weeknights			
Percent (%	Male	Female	Total	
Less than 7 hours	42.0	42.0	42.2	
7 to 9 hours	56.1	55.5	55.4	
10 or more hours	2.0	2.4	2.4	

On weekend nights				
Male	Female	Total		
24.5	24.6	24.7		
66.5	66.1	65.9		
9.0	9.3	9.4		

Students reported the following on 3 or more of the last 7 days:

		Felt tired or sleepy during the day		
H	Percent (%)	Male	Female	Total
0 days		7.6	2.9	4.2
1-2 days		27.0	16.1	18.7
3-5 days		40.3	42.0	41.2
6-7 days		25.0	39.0	35.9

Got enough sleep so that they			
felt rested			
	Male	Female	Total
	17.2	22.4	21.3
	31.9	40.3	37.9
	37.3	28.6	30.7

8.7

10.0

13.6

## **Demographics and Sample Characteristics**

### ■ Age

9.		
18 - 20 years:		60.5 %
21 - 24 years:		30.2 %
25 - 29 years:		4.4 %
30+ years:		4.8 %
Mean age:	21.1 years	
Median age:	20.0 years	
-	-	

#### Gender\*

Stratt	
Female:	70.1 %
Male:	25.9 %
Non-binary:	3.8 %
* See note on page 2 regarding gender categories	

Student status

1st year undergraduate:	26.6 %
2nd year undergraduate:	23.5 %
3rd year undergraduate:	23.8 %
4th year undergraduate:	19.5 %
5th year or more undergraduate:	6.7 %
Master's (MA, MS, MFA, MBA, etc.):	0.0 %
Doctorate (PhD, EdD, MD, JD, etc.):	0.0 %
Not seeking a degree:	0.0 %
Other:	0.0 %
Full-time student:	94.4 %
Part-time student:	5.3 %
Other student:	0.3 %
■ Visa to work or study in the US:	6.1 %
Relationship status	
Not in a relationship:	54.6 %
In a relationship but not married/partnered:	40.4 %
Married/partnered:	5.0 %
-	0.0 / 0
Primary Source of Health Insurance	
College/university sponsored SHIP plan:	6.4 %
College/university sponsored SHIP plan: Parent or guardian's plan:	6.4 % 74.0 %
College/university sponsored SHIP plan: Parent or guardian's plan: Employer (mine or my spouse/partners):	6.4 % 74.0 % 4.2 %
College/university sponsored SHIP plan: Parent or guardian's plan: Employer (mine or my spouse/partners): Medicaid, Medicare, SCHIP, or VA:	6.4 % 74.0 % 4.2 % 7.6 %
College/university sponsored SHIP plan: Parent or guardian's plan: Employer (mine or my spouse/partners): Medicaid, Medicare, SCHIP, or VA: Bought a plan on my own:	6.4 % 74.0 % 4.2 % 7.6 % 1.8 %
College/university sponsored SHIP plan: Parent or guardian's plan: Employer (mine or my spouse/partners): Medicaid, Medicare, SCHIP, or VA: Bought a plan on my own: Don't have health insurance:	6.4 % 74.0 % 4.2 % 7.6 % 1.8 % 3.5 %
College/university sponsored SHIP plan: Parent or guardian's plan: Employer (mine or my spouse/partners): Medicaid, Medicare, SCHIP, or VA: Bought a plan on my own: Don't have health insurance: Don't know if I have health insurance:	6.4 % 74.0 % 4.2 % 7.6 % 1.8 % 3.5 % 0.8 %
College/university sponsored SHIP plan: Parent or guardian's plan: Employer (mine or my spouse/partners): Medicaid, Medicare, SCHIP, or VA: Bought a plan on my own: Don't have health insurance:	6.4 % 74.0 % 4.2 % 7.6 % 1.8 % 3.5 %

### Parent or primary responsibility for someone

else's child/children under [	18 years old:	4.6 %
-------------------------------	---------------	-------

#### Students describe themselves as

Straight/Heterosexual:	77.9 %
Asexual:	1.1 %
Bisexual:	11.1 %
Gay:	1.7 %
Lesbian:	1.6 %
Pansexual:	2.1 %
Queer:	1.3 %
Questioning:	2.8 %
Identity not listed above:	0.4 %

#### Housing

South Asian:

Other Asian:

Campus or university housing:	33.6 %
Fraternity or sorority residence:	2.4 %
Parent/guardian/other family:	28.1 %
Off-campus:	34.4 %
Temporary or "couch surfing":	0.5 %
Don't have a place to live:	0.0 %
Other:	0.9 %

#### Students describe themselves as

American Indian or Native Alaskan	3.0 %
Asian or Asian American	10.0 %
Black or African American	8.1 %
Hispanic or Latino/a/x	9.3 %
Middle Eastern/North African (MENA)	
or Arab Origin:	1.2 %
Native Hawaiian or Other	
Pacific Islander Native:	0.5 %
White:	73.8 %
Biracial or Multiracial:	3.9 %
Identity not listed above:	1.4 %
If Hispanic or Latino/a/x, are you	
Mexican, Mexican American, Chicano:	42.1 %
Puerto Rican:	14.0 %
Puerto Rican: Cuban:	14.0 % 4.6 %
Cuban:	
Cuban: Another Hispanic, Latino/a/x, or	4.6 %
Cuban: Another Hispanic, Latino/a/x, or Spanish Origin:	4.6 %

26.3 % 1.7 %

• First generation students (Proportion of students for whom no parent/guardian have completed a bachelor's degree)	38.4 %
Do you have any of the following?	
Attention Deficit/Hyperactivity Disorder	
(ADD or ADHD):	11.8 %
Autism Spectrum Disorder:	1.6 %
Deaf/Hearing loss:	2.2 %
Learning disability:	4.3 %
Mobility/Dexterity disability:	1.0 %

4.2 %

1.0 %

Blind/low vision:

Speech or language disorder:

### Participated in organized college athletics:

Varsity:	6.1 %
Club sports:	6.6 %
Intramurals:	11.0 %

## ■ Member of a <u>social</u> fraternity or sorority:

Greek member: 11.5 %

### **Demographics of Participating Institutions**

Twenty three postsecondary institutions self-selected to participate in the Fall 2020 ACHA National College Health Assessment and 14,414 surveys were completed by students on these campuses. For the purpose of forming the Reference Group, only institutions located in the United States that surveyed all students or used a random sampling technique, and used the ACHA-NCHA III (currently only offered as a web survey) are included in the analysis, yielding a final data set consisting of 13,373 students at 22 schools. This report includes only data from 8,851

**UNDERGRADUATE students at 22 schools.** Demographic characteristics of the 22 campuses follow.

Demographical Characteristics of the 22 US Postsecondary Institutions Included in the Fall 2020 ACHA-NCHA III UNDERGRADUATE Reference Group		
Campus Characteristic	п	
Type of Institution		
Public	15	
Private	7	
2-year	1	
4-year or above	21	
Location of Campus		
Northeast (CT, ME, MA, NH, NJ, NY, PA, RI, VT)	7	
Midwest (IL, IN, IA, KS, MI, MN, MO, NE, ND, OH, SD, WI)	5	
South (AL, AR, DE, DC, FL, GA, KY, LA, MD, MS, NC, OK, SC,		
TN, TX, VA, WV)	9	
West (AK, AZ, CA, CO, HI, ID, MT, NV, NM, OR, UT, WA, WY)	1	
Campus Size		
< 2,500 students	6	
2,500 – 4,999 students	2	
5,000 – 9,999 students	3	
10,000 – 19,999 students	5	
20,000 students or more	6	
Campus Setting		
Very large city (population over 500,000)	4	
Large city (population 250,000-499,999)	1	
Small city (population 50,000-249,999)	7	
Large town (population 10,000-49,999)	8	
Small town (population 2,500-9,999)	2	
Rural community (population under 2,500)	0	
Carnegie Classification		
Associates Colleges	1	
Baccalaureate Colleges	4	
Baccalaureate/Associates Colleges	0	
Masters Colleges and Universities	5	
Doctoral Universities	9	
Special Focus Institutions	3	
Miscellaneous/Not Classified	0	

Demographical Characteristics of the 22 US Postsecondary Institutions Included in the Fall 2020 ACHA-NCHA III UNDERGRADUATE Reference Group		
Campus Characteristic	n	
ACHA Membership Status		
Institutional Member	18	
Nonmember	4	
Religious Affiliation		
No	18	
Yes	4	
<u>If yes:</u>		
Catholic	2	
Protestant or Other Christian	2	
Postsecondary Minority Institution (US Department of Education)		
No	20	
Yes	2	
<u>*If yes:</u>		
Postsecondary Minority Institution	0	
Historically Black College or University (HBCU)	0	
High Hispanic Enrollment	0	
Hispanic-serving Institution (HSI)	2	
Tribal College or University	0	
Predominately Black Institution	0	
Asian American and Native American Pacific Islander-serving	0	
Alaska Native-serving or Native Hawaiian-serving Institution	0	
Native American-serving Nontribal Institution	0	
*institutions may hold more than one type of minority status		

UNDERGRADUATE STUDENTS	Fall 2020 Institutions
Number of institutions	22
Number of students	8,851
Mean response proportion*	14%
Median response proportion*	11%

\*Calculation of response proportion is based all students surveyed at the institution, and not limited to only undergraduate student responses.